

Near extreme eigenvalues of large random Gaussian matrices and applications

Grégory Schehr
LPTMS, CNRS–Université Paris–Sud XI

Groupe de Travail «Matrices et graphes aléatoires»
Paris (IHP), January 16, 2015

Collaborators:

- Anthony Perret (LPTMS, Orsay)
- Yan V. Fyodorov (Queen Mary, London)

A. Perret, G. S., J. Stat. Phys. **156**, pp. 843–876 (2014)

A. Perret, Y. V. Fyodorov, G. S., in preparation

Extreme value statistics

• Statement of the problem

X_1, X_2, \dots, X_N : N random variables, $P_{\text{joint}}(X_1, X_2, \dots, X_N)$

$$X_{\max} = \max_{1 \leq i \leq N} X_i, F_N(M) = \mathbb{P}(X_{\max} \leq M) \quad \text{Q: } N \rightarrow \infty \quad ?$$

• Fully understood for i.i.d. random variables

Three different universality classes depending on the pdf of X_i : Gumbel, Fréchet, Weibull

Extreme value statistics

• Statement of the problem

$X_1, X_2, \dots, X_N : N$ random variables, $P_{\text{joint}}(X_1, X_2, \dots, X_N)$

$$X_{\max} = \max_{1 \leq i \leq N} X_i, F_N(M) = \mathbb{P}(X_{\max} \leq M) \quad \text{Q: } N \rightarrow \infty \quad ?$$

• Very few exact results for strongly correlated variables


✓ Random walks

✓ Random matrices

• X_{\max} is interesting BUT concerns a single variable among N

Statistics of Near-Extremes

Statistical Physics




see also: ✓ Branching Brownian motion

✓ $1/f$ noise

✓ Brownian motion

✓ ...

Natural sciences (e.g. seismology)



Crowding near the extremes

Statistics of Near-Extremes

How to quantify the crowding close to extreme values ?

- Look at (higher) order statistics: k^{th} maximum

$$X_{\max} = M_{1,N} > M_{2,N} > \cdots > M_{N,N} = X_{\min}$$

and in particular the spacings (gaps) $d_{k,N} = M_{k,N} - M_{k+1,N}$

- Consider the density of near-extremes [Sabhapandit, Majumdar '07](#)

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{i=1, i \neq \text{imax}}^N \overline{\delta(X_{\max} - X_i - r)}$$

Near-extreme eigenvalues of random matrices

- Let J be a real symmetric (or complex Hermitian) $N \times N$ random matrix
- The matrix J has N real eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_N$ which are strongly correlated
- Largest eigenvalue $\lambda_{\max} = \max\{\lambda_1, \lambda_2, \dots, \lambda_N\}$

- Density of near extreme eigenvalues

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\max}}}^N \overline{\delta(\lambda_{\max} - \lambda_i - r)}$$

- A related quantity is the gap between the two largest eigenvalues

see also Witte, Bornemann, Forrester '13

Application: minimizing a quadratic form on the sphere

- Quadratic form on the N -dimensional sphere S_N

$$H[\vec{s}] = -\frac{1}{2} \sum_{i,j=1}^N J_{ij} s_i s_j, \quad (\vec{s})^2 = \sum_{i=1}^N s_i^2 = N$$

$\lambda_1, \lambda_2, \dots, \lambda_N$ are the eigenvalues of J with $\lambda_{\max} = \max_{1 \leq i \leq N} \lambda_i$

- Minimisation of this quadratic form on the sphere

\Rightarrow introduce a Lagrange multiplier z

$$\tilde{H}[\vec{s}, z] = -\frac{1}{2} \sum_{i,j=1}^N J_{ij} s_i s_j + z \left(\sum_{i=1}^N s_i^2 - N \right)$$

$$\min_{\vec{s}, z} \tilde{H}[\vec{s}, z] = \tilde{H}[\vec{s}_{\max}, z_{\max}] = -N \frac{\lambda_{\max}}{2} \text{ with } \begin{cases} J \vec{s}_{\max} = \lambda_{\max} \vec{s}_{\max} \\ z_{\max} = \frac{\lambda_{\max}}{2} \end{cases}$$

Application: minimizing a quadratic form on the sphere

$$\tilde{H}[\vec{s}, z] = -\frac{1}{2} \sum_{i,j=1}^N J_{ij} s_i s_j + z \left(\sum_{i=1}^N s_i^2 - N \right)$$

$$\min_{\vec{s}, z} \tilde{H}[\vec{s}, z] = \tilde{H}[\vec{s}_{\max}, z_{\max}] = -N \frac{\lambda_{\max}}{2} \text{ with } \begin{cases} J \vec{s}_{\max} = \lambda_{\max} \vec{s}_{\max} \\ z_{\max} = \frac{\lambda_{\max}}{2} \end{cases}$$

• Eigenvalues of the Hessian matrix at the minimum \vec{s}_{\max}, z_{\max}

spectrum of $\left. \frac{\delta^2 \tilde{H}}{\delta s_i \delta s_j} \right|_{\vec{s}_{\max}, z_{\max}}$ is $\{0, \lambda_{\max} - \lambda_1, \lambda_{\max} - \lambda_2, \dots, \lambda_{\max} - \lambda_N\}$

Reminding that $\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\max}}}^N \overline{\delta(\lambda_{\max} - \lambda_i - r)}$

$\Rightarrow \rho_{\text{DOS}}(r, N)$ is the mean eigenvalue density of $\left. \frac{\delta^2 \tilde{H}}{\delta s_i \delta s_j} \right|_{\vec{s}_{\max}, z_{\max}}$

Application to spherical fully connected spin-glasses

- Dynamics of the spherical fully connected spin-glass model
(spherical Sherrington-Kirkpatrick model)

Cugliandolo, Dean '95

Ben Arous, Dembo, Guionnet '06

$$\frac{\partial s_i(t)}{\partial t} = -\frac{\delta}{\delta s_i(t)} \tilde{H}[\vec{s}(t), z(t)] + h_i(t) + \zeta_i(t) \quad \leftarrow \text{temp. } T$$

where $\tilde{H}[\vec{s}, z] = -\frac{1}{2} \sum_{i,j=1}^N J_{ij} s_i s_j + z \left(\sum_{i=1}^N s_i^2 - N \right)$ infinitesimal
mag. field

and J belongs to the GOE ensemble of RMT with variance $\overline{J_{ij}^2} = \frac{1}{N}$

- Random initial condition at time $t = 0$: $\vec{s}(t = 0) = \sum_{\alpha=1}^N \vec{s}_{\alpha}$

normalized eigenvectors of J : $\vec{s}_{\alpha} \mid J \vec{s}_{\alpha} = \lambda_{\alpha} \vec{s}_{\alpha}$

- Relaxational dynamics characterized by **two-time quantities**

Application to spherical fully connected spin-glasses

$$\frac{\partial s_i(t)}{\partial t} = \sum_{j=1}^N J_{ij} s_j(t) - z(t) s_i(t) + h_i(t), \quad T = 0$$

- Relaxational dynamics characterized by **two-time quantities** $t > t'$

Correlation function

$$C(t, t') = \frac{1}{N} \sum_{i=1}^N \overline{s_i(t) s_i(t')}$$

Response function

$$R(t, t') = \frac{1}{N} \sum_{i=1}^N \left. \overline{\frac{\delta s_i(t)}{\delta h_i(t')}} \right|_{h=0}$$

- In the quasi-stationary regime, for $t, t' \gg 1$, $t - t' = \tau$ fixed


$$R(t, t') \sim \int_0^\infty e^{-r(t-t')} \rho_{\text{DOS}}(r, N) dr$$

Perret, Fyodorov, G. S. '14

see also Kurchan, Laloux '96

Near-extreme eigenvalues of random matrices

$$P_{\text{joint}}(\lambda_1, \lambda_2, \dots, \lambda_N) = \frac{1}{Z_N} \prod_{i < j} |\lambda_i - \lambda_j|^\beta e^{-N \frac{\beta}{2} \sum_{i=1}^N \lambda_i^2}$$



✓ Density of near extreme eigenvalues

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\max}}}^N \overline{\delta(\lambda_{\max} - \lambda_i - r)}$$

✓ 1st gap

$$\rho_{\text{GAP}}(r, N) dr = \mathbb{P}[(\Lambda_{1,N} - \Lambda_{2,N}) \in [r, r + dr]]$$

Density of near extreme eigenvalues in GUE

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\text{max}}}}^N \overline{\delta(\lambda_{\text{max}} - \lambda_i - r)}$$

- Fluctuations of the largest eigenvalue $\lambda_{\text{max}} = \max_{1 \leq i \leq N} \lambda_i$

$$\lambda_{\text{max}} = \sqrt{2} + N^{-2/3} \underbrace{\frac{1}{\sqrt{2}} \chi_{\beta}}$$

Tracy-Widom

- Depending on (r, N) one expects two different regimes

$$\checkmark r = O(N^0) \quad \text{bulk regime}$$


$$\checkmark r = O(N^{-2/3}) \quad \text{edge regime}$$

A detour by the density of eigenvalues of random matrices


$$\rho(\lambda, N) = \frac{1}{N} \sum_{i=1}^N \overline{\delta(\lambda - \lambda_i)}$$

Two regimes: bulk and edge regime Bowick & Brézin '91 , Forrester '93

$$\rho(\lambda, N) \sim \begin{cases} \rho_{\text{bulk}}(\lambda) , & \lambda = O(N^0) \text{ \& } |\lambda| < \sqrt{2} \\ \sqrt{2}N^{-1/3} \rho_{\text{edge}}[(\lambda - \sqrt{2})\sqrt{2}N^{2/3}] , & |\lambda - \sqrt{2}| = O(N^{-2/3}) \end{cases}$$



$$\rho_{\text{bulk}}(x) = \frac{1}{\pi} \sqrt{2 - x^2}$$



$$\rho_{\text{edge}}(x) = [\text{Ai}'(x)]^2 - x \text{Ai}^2(x)$$

A detour by the density of eigenvalues of GUE random matrices

$$\rho(\lambda, N) = \frac{1}{N} \sum_{i=1}^N \overline{\delta(\lambda - \lambda_i)}$$

- Two regimes: bulk and edge regime

$$\rho(\lambda, N) \sim \begin{cases} \rho_{\text{bulk}}(\lambda) , \lambda = O(N^0) \ \& \ |\lambda| < \sqrt{2} \\ \sqrt{2}N^{-1/3} \rho_{\text{edge}}[(\lambda - \sqrt{2})\sqrt{2}N^{2/3}] , \ |\lambda - \sqrt{2}| = O(N^{-2/3}) \end{cases}$$

- **Matching** between the bulk and edge regimes

$$\rho_{\text{edge}}(x) \sim \begin{cases} \frac{1}{\pi} \sqrt{-x} , \ x \rightarrow -\infty & \text{matching with Wigner semi-circle} \\ e^{-\frac{2\beta}{3}x^{3/2}} , \ x \rightarrow +\infty & \text{coincides with the right tail of TW} \end{cases}$$

Density of near extreme eigenvalues: results

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\text{max}}}}^N \overline{\delta(\lambda_{\text{max}} - \lambda_i - r)}$$

$$\rho_{\text{DOS}}(r, N) \sim \begin{cases} \tilde{\rho}_{\text{bulk}}(r) , & r = O(N^0) \text{ \& } 0 < r < 2\sqrt{2} \\ \sqrt{2}N^{-1/3} \tilde{\rho}_{\text{edge}}(r\sqrt{2}N^{2/3}) , & r = O(N^{-2/3}) \end{cases}$$

A. Perret, G. S. '14

• In the bulk $r \sim O(N^0)$:

✓ $\rho_{\text{DOS}}(r, N)$ is insensitive to the fluctuations of $\lambda_{\text{max}} \sim \sqrt{2}$

$$\checkmark \rho_{\text{DOS}}(r, N) \sim \frac{1}{N} \sum_{i=1}^N \overline{\delta(\sqrt{2} - \lambda_i - r)} = \rho(\sqrt{2} - r, N)$$

$$\Rightarrow \checkmark \tilde{\rho}_{\text{bulk}}(x) = \frac{1}{\pi} \sqrt{x(2\sqrt{2} - x)} \text{ shifted Wigner semi-circle}$$

Density of near extreme eigenvalues: results

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\text{max}}}}^N \overline{\delta(\lambda_{\text{max}} - \lambda_i - r)}$$

$$\rho_{\text{DOS}}(r, N) \sim \begin{cases} \tilde{\rho}_{\text{bulk}}(r) , & r = O(N^0) \text{ \& } 0 < r < 2\sqrt{2} \\ \sqrt{2}N^{-1/3} \tilde{\rho}_{\text{edge}}(r\sqrt{2}N^{2/3}) , & r = O(N^{-2/3}) \end{cases}$$

• At the edge $r \sim O(N^{-2/3})$, a non trivial function

$$\tilde{\rho}_{\text{edge}}(x) \sim \begin{cases} a_{\beta} x^{\beta} , & x \rightarrow 0 \\ \frac{\sqrt{x}}{\pi} , & x \rightarrow \infty \end{cases}$$

Consequences on the dynamics of the spherical fully connected spin-glass model

$$\frac{\partial s_i(t)}{\partial t} = \sum_{j=1}^N J_{ij} s_j(t) - z(t) s_i(t) + h_i(t), \quad T = 0$$

- In the quasi-stationary regime, for $t, t' \gg 1$, $t - t' = \tau$

$$R(t, t') \sim \int_0^\infty e^{-r(t-t')} \rho_{\text{DOS}}(r, N) dr$$

two regimes: $\rho_{\text{DOS}}(r, N) \sim \begin{cases} \tilde{\rho}_{\text{bulk}}(r), & r = O(N^0) \text{ \& } 0 < r < 2\sqrt{2} \\ \sqrt{2}N^{-1/3} \tilde{\rho}_{\text{edge}}(r\sqrt{2}N^{2/3}), & r = O(N^{-2/3}) \end{cases}$

- Two temporal regimes for the response function

✓ $1 \ll t, t' \ll N^{2/3}$

Cugliandolo, Dean '95
Ben Arous, Dembo, Guionnet '06

✓ $t, t' = O(N^{2/3})$

Perret, Fyodorov, G. S. '15

Consequences on the dynamics of the spherical fully connected spin-glass model

$$R(t, t') \sim \int_0^\infty e^{-r(t-t')} \rho_{\text{DOS}}(r, N) dr$$

• For $\tau = (t - t') \sim \mathcal{O}(N^{2/3})$

$$R(t, t') \sim \frac{1}{N} f_R \left(\frac{t - t'}{N^{2/3}} \right), \quad f_R(x) = \int_0^\infty e^{-\frac{r}{\sqrt{2}}x} \tilde{\rho}_{\text{edge}}(x) dx$$

universal !

• Asymptotic behaviors

$$f_R(x) \sim \begin{cases} x^{-3/2}, & x \rightarrow 0 \\ x^{-2}, & x \rightarrow \infty \end{cases}$$

• Can one compute the full universal function $\tilde{\rho}_{\text{edge}}(x)$?

\implies an exact calculation for GUE

Perret, G. S. '14

Density of near extreme eigenvalues for GUE

Perret, G. S. '14

$$\tilde{\rho}_{\text{edge}}(\tilde{r}) = \frac{2^{1/3}}{\pi} \int_{-\infty}^{\infty} \left(\tilde{f}^2(\tilde{r}, x) - \left(\int_x^{\infty} q(u) \tilde{f}(\tilde{r}, u) du \right)^2 \right) \mathcal{F}_2(x) dx$$

$$\mathcal{F}_2(x) = \exp \left[- \int_x^{\infty} (u - x) q^2(u) du \right] , \quad \begin{cases} q'' = 2q^3 + xq , \\ q(x) \sim \text{Ai}(x) \text{ for } x \rightarrow \infty , \end{cases}$$

$$\partial_x^2 \tilde{f}(\tilde{r}, x) - [x + 2q^2(x)] \tilde{f}(\tilde{r}, x) = -\tilde{r} \tilde{f}(\tilde{r}, x) , \quad \tilde{f}(\tilde{r}, x) \underset{x \rightarrow \infty}{\sim} 2^{-1/6} \sqrt{\pi} \text{Ai}(x - \tilde{r})$$

$\tilde{f}(\tilde{r}, x)$ is related to a solution of the Lax pair associated to Painlevé XXXIV

Density of near extreme eigenvalues for GUE

$$\tilde{\rho}_{\text{edge}}(\tilde{r}) = \frac{2^{1/3}}{\pi} \int_{-\infty}^{\infty} \left(\tilde{f}^2(\tilde{r}, x) - \left(\int_x^{\infty} q(u) \tilde{f}(\tilde{r}, u) du \right)^2 \right) \mathcal{F}_2(x) dx$$

$$\partial_x^2 \tilde{f}(\tilde{r}, x) - [x + 2q^2(x)] \tilde{f}(\tilde{r}, x) = -\tilde{r} \tilde{f}(\tilde{r}, x), \quad \tilde{f}(\tilde{r}, x) \underset{x \rightarrow \infty}{\sim} 2^{-1/6} \sqrt{\pi} \text{Ai}(x - \tilde{r})$$

$\tilde{f}(\tilde{r}, x)$ is related to a solution of the Lax pair associated to Painlevé XXXIV

$$\frac{\partial}{\partial \tilde{r}} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix} = \tilde{\mathbf{A}} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix}, \quad \frac{\partial}{\partial x} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix} = \tilde{\mathbf{B}} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix}$$

$$\tilde{\mathbf{A}} = \begin{pmatrix} -\frac{q'(x)}{q(x)} & 1 + \frac{q^2(x)}{\tilde{r}} \\ -\tilde{r} - \frac{\int_x^{\infty} q^2(u) du}{q^2(x)} & \frac{q'(x)}{q(x)} \end{pmatrix}, \quad \tilde{\mathbf{B}} = \begin{pmatrix} \frac{q'(x)}{q(x)} & -1 \\ \tilde{r} & -\frac{q'(x)}{q(x)} \end{pmatrix}$$

$$\tilde{f}(\tilde{r}, x) \underset{\tilde{r} \rightarrow \infty}{\sim} 2^{-1/6} \tilde{r}^{-1/4} \sin \left(\frac{2}{3} \tilde{r}^{3/2} - x \sqrt{\tilde{r}} + \frac{\pi}{4} \right) + \mathcal{O}(\tilde{r}^{-3/4}),$$


$$\tilde{g}(\tilde{r}, x) \underset{\tilde{r} \rightarrow \infty}{\sim} 2^{-1/6} \tilde{r}^{1/4} \cos \left(\frac{2}{3} \tilde{r}^{3/2} - x \sqrt{\tilde{r}} + \frac{\pi}{4} \right) + \mathcal{O}(\tilde{r}^{-1/4})$$

Density of near extreme eigenvalues: results

$$\tilde{\rho}_{\text{edge}}(\tilde{r}) = \frac{2^{1/3}}{\pi} \int_{-\infty}^{\infty} \left(\tilde{f}^2(\tilde{r}, x) - \left(\int_x^{\infty} q(u) \tilde{f}(\tilde{r}, u) du \right)^2 \right) \mathcal{F}_2(x) dx$$

Asymptotic behaviors

$$\tilde{\rho}_{\text{edge}}(\tilde{r}) \sim \begin{cases} \frac{1}{2} \tilde{r}^2 + a_4 \tilde{r}^4 + \mathcal{O}(\tilde{r}^6) & , \tilde{r} \rightarrow 0 \\ \frac{\sqrt{\tilde{r}}}{\pi} & , \tilde{r} \rightarrow \infty \end{cases}$$



Outline


- Exact formulas for $\rho_{\text{DOS}}(r, N)$ & $\rho_{\text{GAP}}(r, N)$ for finite N
- Asymptotic analysis for large N
- Comparison with existing results
- Conclusion and related open problems

Outline

- Exact formulas for $\rho_{\text{DOS}}(r, N)$ & $\rho_{\text{GAP}}(r, N)$ for finite N
- Asymptotic analysis for large N
- Comparison with existing results
- Conclusion and related open problems

An exact formula for $\rho_{\text{DOS}}(r, N)$

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\text{max}}}}^N \overline{\delta(\lambda_{\text{max}} - \lambda_i - r)}$$



$$\rho_{\text{DOS}}(r, N) = N \int_{-\infty}^{\infty} dy \underbrace{\int_{-\infty}^y d\lambda_1 \int_{-\infty}^y d\lambda_2 \cdots \int_{-\infty}^y d\lambda_{N-2} P_{\text{joint}}(\lambda_1, \lambda_2, \dots, \lambda_{N-2}, y-r, y)}_{\text{two-point correlations for conditioned eigenvalues}}$$

two-point correlations for **conditioned** eigenvalues

• After some manipulations one obtains


$$\rho_{\text{DOS}}(r, N) = \frac{N(N-2)!}{Z_N} \int_{-\infty}^{\infty} dy \prod_{k=0}^{N-1} h_k(y) \begin{vmatrix} K_N(y-r, y-r) & K_N(y-r, y) \\ K_N(y, y-r) & K_N(y, y) \end{vmatrix}$$

with the kernel

$$K_N(\lambda, \lambda') = \sum_{k=0}^{N-1} \frac{1}{h_k(y)} \pi_k(\lambda, y) \pi_k(\lambda', y) e^{-\frac{\lambda^2 + \lambda'^2}{2}}$$

An exact formula for $\rho_{\text{DOS}}(r, N)$

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\text{max}}}}^N \overline{\delta(\lambda_{\text{max}} - \lambda_i - r)}$$




$$\rho_{\text{DOS}}(r, N) = \frac{N(N-2)!}{Z_N} \int_{-\infty}^{\infty} dy \underbrace{\prod_{k=0}^{N-1} h_k(y)}_{\propto F_N(y) = \mathbb{P}(\lambda_{\text{max}} \leq y)} \underbrace{\begin{vmatrix} K_N(y-r, y-r) & K_N(y-r, y) \\ K_N(y, y-r) & K_N(y, y) \end{vmatrix}}_{\text{2-point correlations}}$$

Finally a useful formula is

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \int_{-\infty}^{\infty} dy [F'_N(y) K_N(y-r, y-r) - F_N(y) K_N^2(y, y-r)]$$

An exact formula for the PDF of the first gap

$$\rho_{\text{GAP}}(r, N)dr = \mathbb{P}[(\Lambda_{1,N} - \Lambda_{2,N}) \in [r, r + dr]]$$



$$\rho_{\text{GAP}}(r, N) = N(N-1) \int_{-\infty}^{+\infty} dy \int_{-\infty}^y d\lambda_1 \int_{-\infty}^y d\lambda_2 \cdots \int_{-\infty}^y d\lambda_{N-2} P_{\text{joint}}(\lambda_1, \dots, \lambda_{N-2}, y, y+r)$$

Reminding that

$$\rho_{\text{DOS}}(r, N) = N \int_{-\infty}^{\infty} dy \int_{-\infty}^y d\lambda_1 \int_{-\infty}^y d\lambda_2 \cdots \int_{-\infty}^y d\lambda_{N-2} P_{\text{joint}}(\lambda_1, \lambda_2, \dots, \lambda_{N-2}, y-r, y)$$

one gets

$$\rho_{\text{GAP}}(r, N) = (N-1)\rho_{\text{DOS}}(-r, N)$$

Outline

- Exact formulas for $\rho_{\text{DOS}}(r, N)$ & $\rho_{\text{GAP}}(r, N)$ for finite N
- Orthogonal polynomials (OPs) on the semi-infinite real line
- Comparison with existing results
- Conclusion and related open problems

Orthogonal polynomials

$$P_{\text{joint}}(\lambda_1, \lambda_2, \dots, \lambda_N) = \frac{1}{Z_N} \prod_{i < j} (\lambda_i - \lambda_j)^2 e^{-\sum_{i=1}^N \lambda_i^2}$$

$$\lambda_{\text{max}} = \max_{1 \leq i \leq N} \lambda_i$$

- Cumulative distribution function of λ_{max}

$$F_N(y) = \text{Pr.}(\max_{1 \leq i \leq N} \lambda_i \leq y)$$

$$F_N(y) = \frac{1}{Z_N} \int_{-\infty}^y d\lambda_1 \cdots \int_{-\infty}^y d\lambda_N \prod_{i < j} (\lambda_i - \lambda_j)^2 e^{-\sum_{i=1}^N \lambda_i^2}$$

- Orthogonal polynomials on the semi-infinite real line

$$\langle \pi_k, \pi_{k'} \rangle = \int_{-\infty}^y d\lambda \pi_k(\lambda, y) \pi_{k'}(\lambda, y) e^{-\lambda^2} = \delta_{k,k'} h_k(y)$$

$$\pi_k(\lambda, y) = \lambda^k + \dots$$

$$F_N(y) = \frac{N!}{Z_N} \prod_{j=0}^{N-1} h_j(y)$$

C. Nadal, S. N. Majumdar '13

Physical picture associated to the OP sytem


$$F_N(y) = \Pr.(\max_{1 \leq i \leq N} \lambda_i \leq y)$$

$$F_N(y) = \frac{1}{Z_N} \int_{-\infty}^y d\lambda_1 \cdots \int_{-\infty}^y d\lambda_N \prod_{i < j} (\lambda_i - \lambda_j)^2 e^{-\sum_{i=1}^N \lambda_i^2}$$

- Coulomb gas with a **wall** located in y

for a review see S. N. Majumdar, G. S. '13

- Moving the wall through the edge: the density $\rho_y(\lambda)$



described by a double scaling limit

see also T. Claeys, A. Kuijlaars '08, «...when the soft edge meets the hard edge»

Large N analysis of $\rho_{\text{DOS}}(N, r)$

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\text{max}}}}^N \langle \delta(\lambda_{\text{max}} - \lambda_i - r) \rangle$$

• Edge regime : $r = \mathcal{O}(N^{-\frac{1}{6}})$

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \int_{-\infty}^{\infty} dy [F'_N(y) K_N(y-r, y-r) - F_N(y) K_N^2(y, y-r)]$$

• Analysis of the kernel in the double scaling limit

$$K_N(\lambda, \lambda') = \sum_{k=0}^{N-1} \psi_k(\lambda, y) \psi_k(\lambda', y) \quad \text{where} \quad \psi_k(\lambda, y) = \frac{1}{\sqrt{h_k(y)}} \pi_k(\lambda, y) e^{-\frac{\lambda^2}{2}}$$

Find a solution of the recurrence in the double scaling limit

$$\lambda \psi_N(\lambda, y) = \sqrt{R_{N+1}(y)} \psi_{N+1}(\lambda, y) + S_N(y) \psi_N(\lambda, y) + \sqrt{R_N(y)} \psi_{N-1}(\lambda, y)$$

Large N analysis of $\rho_{\text{DOS}}(N, r)$

$$\rho_{\text{DOS}}(r, N) = \frac{1}{N-1} \sum_{\substack{i=1 \\ i \neq i_{\text{max}}}}^N \langle \delta(\lambda_{\text{max}} - \lambda_i - r) \rangle$$

$$\lambda \psi_N(\lambda, y) = \sqrt{R_{N+1}(y)} \psi_{N+1}(\lambda, y) + S_N(y) \psi_N(\lambda, y) + \sqrt{R_N(y)} \psi_{N-1}(\lambda, y)$$

• In the double scaling limit, one has

C. Nadal, S. N. Majumdar '13

$$\begin{cases} R_N(y) &= \frac{N}{2} \left(1 - N^{-\frac{2}{3}} q^2(x) + \mathcal{O}(N^{-1}) \right) \\ S_N(y) &= -\frac{N^{-\frac{1}{6}}}{\sqrt{2}} q^2(x) + \mathcal{O}(N^{-\frac{1}{2}}) \end{cases} \text{ with } \begin{cases} q'' = 2q^3 + xq, \\ q(x) \sim \text{Ai}(x) \text{ for } x \rightarrow \infty, \end{cases}$$

$$x = \sqrt{2} N^{\frac{1}{6}} (y - \sqrt{2N})$$

• In the double scaling limit the recurrence relation is solved by

$$\psi_N(y - r, y) = \frac{2^{1/4}}{\sqrt{\pi}} N^{-\frac{1}{12}} G \left(\sqrt{2} N^{1/6} r, \sqrt{2} N^{1/6} (y - \sqrt{2N}) \right) + \mathcal{O}(N^{-\frac{5}{12}})$$

$$-\partial_x^2 G(\tilde{r}, x) + [x + 2q^2(x)] G(\tilde{r}, x) = \tilde{r} G(\tilde{r}, x)$$

Large N analysis of $\rho_{\text{DOS}}(N, r)$

- After some more computations...

$$K_N \left(y - \frac{\tilde{r}}{\sqrt{2}} N^{-1/6}, y - \frac{\tilde{r}'}{\sqrt{2}} N^{-1/6} \right) \underset{N \rightarrow \infty}{\sim} N^{1/6} 2^{5/6} \frac{\tilde{f}(\tilde{r}, x) \tilde{g}(\tilde{r}', x) - \tilde{f}(\tilde{r}', x) \tilde{g}(\tilde{r}, x)}{\pi(\tilde{r} - \tilde{r}')}$$

where $\tilde{f}(\tilde{r}, x)$, $\tilde{g}(\tilde{r}, x)$ solve the Lax pair for Painlevé XXXIV

$$\frac{\partial}{\partial \tilde{r}} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix} = \tilde{\mathbf{A}} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix}, \quad \frac{\partial}{\partial x} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix} = \tilde{\mathbf{B}} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix}$$

$$\tilde{\mathbf{A}} = \begin{pmatrix} -\frac{q'(x)}{q(x)} & 1 + \frac{q^2(x)}{\tilde{r}} \\ -\tilde{r} - \frac{\int_x^\infty q^2(u) du}{q^2(x)} & \frac{q'(x)}{q(x)} \end{pmatrix}, \quad \tilde{\mathbf{B}} = \begin{pmatrix} \frac{q'(x)}{q(x)} & -1 \\ \tilde{r} & -\frac{q'(x)}{q(x)} \end{pmatrix}$$

$$\tilde{f}(\tilde{r}, x) \underset{\tilde{r} \rightarrow \infty}{\sim} 2^{-1/6} \tilde{r}^{-1/4} \sin \left(\frac{2}{3} \tilde{r}^{3/2} - x\sqrt{\tilde{r}} + \frac{\pi}{4} \right) + \mathcal{O}(\tilde{r}^{-3/4}),$$

$$\tilde{g}(\tilde{r}, x) \underset{\tilde{r} \rightarrow \infty}{\sim} 2^{-1/6} \tilde{r}^{1/4} \cos \left(\frac{2}{3} \tilde{r}^{3/2} - x\sqrt{\tilde{r}} + \frac{\pi}{4} \right) + \mathcal{O}(\tilde{r}^{-1/4})$$

see T. Claeys, A. Kuijlaars '07 for a (rigorous) derivation using RH

Density of near extreme eigenvalues: results

$$\tilde{\rho}_{\text{edge}}(\tilde{r}) = \frac{2^{1/3}}{\pi} \int_{-\infty}^{\infty} \left(\tilde{f}^2(\tilde{r}, x) - \left(\int_x^{\infty} q(u) \tilde{f}(\tilde{r}, u) du \right)^2 \right) \mathcal{F}_2(x) dx$$

$$\partial_x^2 \tilde{f}(\tilde{r}, x) - [x + 2q^2(x)] \tilde{f}(\tilde{r}, x) = -\tilde{r} \tilde{f}(\tilde{r}, x), \quad \tilde{f}(\tilde{r}, x) \underset{x \rightarrow \infty}{\sim} 2^{-1/6} \sqrt{\pi} \text{Ai}(x - \tilde{r})$$

$\tilde{f}(\tilde{r}, x)$ is related to a solution of the Lax pair associated to Painlevé XXXIV

$$\frac{\partial}{\partial \tilde{r}} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix} = \tilde{\mathbf{A}} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix}, \quad \frac{\partial}{\partial x} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix} = \tilde{\mathbf{B}} \begin{pmatrix} \tilde{f}(\tilde{r}, x) \\ \tilde{g}(\tilde{r}, x) \end{pmatrix}$$

$$\tilde{\mathbf{A}} = \begin{pmatrix} -\frac{q'(x)}{q(x)} & 1 + \frac{q^2(x)}{\tilde{r}} \\ -\tilde{r} - \frac{\int_x^{\infty} q^2(u) du}{q^2(x)} & \frac{q'(x)}{q(x)} \end{pmatrix}, \quad \tilde{\mathbf{B}} = \begin{pmatrix} \frac{q'(x)}{q(x)} & -1 \\ \tilde{r} & -\frac{q'(x)}{q(x)} \end{pmatrix}$$

$$\tilde{f}(\tilde{r}, x) \underset{\tilde{r} \rightarrow \infty}{\sim} 2^{-1/6} \tilde{r}^{-1/4} \sin \left(\frac{2}{3} \tilde{r}^{3/2} - x \sqrt{\tilde{r}} + \frac{\pi}{4} \right) + \mathcal{O}(\tilde{r}^{-3/4}),$$

$$\tilde{g}(\tilde{r}, x) \underset{\tilde{r} \rightarrow \infty}{\sim} 2^{-1/6} \tilde{r}^{1/4} \cos \left(\frac{2}{3} \tilde{r}^{3/2} - x \sqrt{\tilde{r}} + \frac{\pi}{4} \right) + \mathcal{O}(\tilde{r}^{-1/4})$$

Typical fluctuations of the gap: results

A. Perret, G. S. '13

using $\rho_{\text{GAP}}(r, N) = (N - 1)\rho_{\text{DOS}}(-r, N)$

we obtain $\rho_{\text{GAP}}(r, N) = \sqrt{2}N^{1/6}\tilde{\rho}_{\text{typ}}(r\sqrt{2}N^{1/6})$

$$\tilde{\rho}_{\text{typ}}(\tilde{r}) = \frac{2^{1/3}}{\pi} \int_{-\infty}^{\infty} \left(\tilde{f}^2(-\tilde{r}, x) - \left(\int_x^{\infty} q(u) \tilde{f}(-\tilde{r}, u) du \right)^2 \right) \mathcal{F}_2(x) dx$$

$$\partial_x^2 \tilde{f}(-\tilde{r}, x) - [x + 2q^2(x)] \tilde{f}(-\tilde{r}, x) = \tilde{r} \tilde{f}(-\tilde{r}, x), \quad \tilde{f}(-\tilde{r}, x) \underset{x \rightarrow \infty}{\sim} 2^{-1/6} \sqrt{\pi} \text{Ai}(x + \tilde{r})$$

from which we obtain the asymptotic behaviors

$$\tilde{\rho}_{\text{typ}}(\tilde{r}) = \begin{cases} \frac{1}{2}\tilde{r}^2 + a_4\tilde{r}^4 + \mathcal{O}(\tilde{r}^6), & \tilde{r} \rightarrow 0 \\ A \exp\left(-\frac{4}{3}\tilde{r}^{3/2} + \frac{8}{3}\sqrt{2}\tilde{r}^{3/4}\right) \tilde{r}^{-21/32} \left(1 - \frac{1405\sqrt{2}}{1536}\tilde{r}^{-3/4} + \mathcal{O}(\tilde{r}^{-3/2})\right), & \tilde{r} \rightarrow +\infty \end{cases}$$

with $A = 2^{-91/48} e^{\zeta'(-1)} / \sqrt{\pi}$

and $a_4 = -0.393575\dots$ complicated integral involving $q(x)$

Outline

- Exact formulas for $\rho_{\text{DOS}}(r, N)$ & $\rho_{\text{GAP}}(r, N)$ for finite N
- Orthogonal polynomials (OPs) on the semi-infinite real line
- Comparison with existing results
- Conclusion and related open problems

Relations with existing results

- Density of near extreme eigenvalues was not studied in RMT
(to my knowledge)
- Previous studies of the PDF of the first gap
 - ✓ an expression in terms of a Fredholm determinant Forrester '93
 - ✓ an expression in terms of Painlevé transcendents
Witte, Bornemann, Forrester '13and a numerical computation of the formula in terms of a Fredholm determinant

PDF of the first gap: the formula of Witte, Bornemann, Forrester

$$\tilde{\rho}_{\text{typ}}(\tilde{r}) = \int_{-\infty}^{\infty} p_{(2)}^{\text{soft}}(t, t - \tilde{r}) dt$$

$$p_{(2)}^{\text{soft}}(t, t - \tilde{r}) = \frac{t^{-5/2}}{4\pi} p_{(1)}^{\text{soft}}(t) \exp\left(-\frac{4}{3}t^{3/2}\right) \exp\left(\int_{2^{1/3}t}^{\infty} dy \left\{ (2q_{3/2} + \frac{4}{p_{3/2}})(-y) - \sqrt{2y} - \frac{5}{2y} \right\}\right) \\ \times (U\partial_x V - V\partial_x U)(-2^{1/3}\tilde{r}; -2^{1/3}t)$$

with

$$p_{(1)}^{\text{soft}}(t) = K^{\text{soft}}(t, t) \exp\left(-\int_s^{\infty} dt \left(\sigma_{\text{II}}(t) - \frac{d}{dt} \log K^{\text{soft}}(t, t)\right)\right) \text{ and } K^{\text{soft}}(x, y) = \frac{\text{Ai}(x)\text{Ai}'(y) - \text{Ai}(y)\text{Ai}'(x)}{x - y}$$

$$\sigma_{\text{II}}(t) = -2^{1/3}H(-2^3t) \quad \text{with} \quad H = -\frac{1}{2} (2q_{\alpha}^2 - p_{\alpha} + t) p_{\alpha} - 2q_{\alpha}, \alpha = 3/2$$


$$\partial_t q_{\alpha} = \partial_{p_{\alpha}} H = p_{\alpha} - q_{\alpha}^2 - \frac{1}{2}t, \quad \partial_t p_{\alpha} = \partial_{q_{\alpha}} H = 2q_{\alpha}p_{\alpha} + 2$$

$$\partial_x \begin{pmatrix} U \\ V \end{pmatrix} = \left\{ \begin{pmatrix} 0 & 0 \\ -\frac{1}{2} & 0 \end{pmatrix} x + \begin{pmatrix} -q_{3/2} - \frac{2}{p_{3/2}} & -1 \\ \frac{1}{2}(t - p_{3/2}) + [q_{3/2} + \frac{2}{p_{3/2}}]^2 & q_{3/2} + \frac{2}{p_{3/2}} \end{pmatrix} + \begin{pmatrix} 1 & p_{3/2} \\ 0 & -1 \end{pmatrix} \frac{1}{x} \right\} \begin{pmatrix} U \\ V \end{pmatrix},$$

$$\partial_t \begin{pmatrix} U \\ V \end{pmatrix} = \left\{ \begin{pmatrix} 0 & 0 \\ \frac{1}{2} & 0 \end{pmatrix} x + \begin{pmatrix} 0 & 1 \\ 0 & -2[q_{3/2} + \frac{2}{p_{3/2}}] \end{pmatrix} \right\} \begin{pmatrix} U \\ V \end{pmatrix}$$

Showing that this formula coincides with ours is still challenging...

PDF of the first gap: the numerical evaluation of Witte, Bornemann, Forrester




Conclusion and related open questions

- Applications to the relaxational dynamics of mean-field spin glass
- Exact results for the statistics of near extreme eigenvalues of GUE
- A new formula for the PDF of the first gap in terms of Painlevé transcendents (precise asymptotics)

$$\tilde{\rho}_{\text{typ}}(\tilde{r}) = \begin{cases} \frac{1}{2}\tilde{r}^2 + a_4\tilde{r}^4 + \mathcal{O}(\tilde{r}^6) , & \tilde{r} \rightarrow 0 \\ A \exp\left(-\frac{4}{3}\tilde{r}^{3/2} + \frac{8}{3}\sqrt{2}\tilde{r}^{3/4}\right) \tilde{r}^{-21/32} \left(1 - \frac{1405\sqrt{2}}{1536}\tilde{r}^{-3/4} + \mathcal{O}(\tilde{r}^{-3/2})\right) , & \tilde{r} \rightarrow +\infty \end{cases}$$

- What about GOE and GSE (skew orthogonal polynomials) ?
- What about Laguerre-Wishart matrices ?



Stochastic processes and random matrices

July 6-31, 2015

<http://lptms.u-psud.fr/workshop/randmat/>

Organizing committee

Alexander Altland

Institut für Theoretische Physik, Universität Köln, Germany

Yan Fyodorov

School of Mathematical Sciences, Queen Mary University of London, Great-Britain

Neil O'Connell

Mathematics Institute, University of Warwick, Great-Britain

Grégory Schehr

Laboratoire de Physique Théorique et Modèles Statistiques, CNRS, Université Paris-Sud, France

Long Courses

Alexei Borodin (Massachusetts Institute of Technology)

Alice Guionnet (Massachusetts Institute of Technology)

Pierre Le Doussal (CNRS, Ecole Normale Supérieure, Paris)

Satya Majumdar (CNRS, Université Paris-Sud, Orsay)

Herbert Spohn (Universität München)

Bálint Virág (University of Toronto)

Integrability and the Kardar-Parisi-Zhang universality class

An introduction to free probability

Replicas, renormalization and integrability in random systems

Recent applications of random matrices in statistical physics

The Kardar-Parisi-Zhang equation – a statistical physics perspective

β -ensembles and random Schrödinger operators

Short Courses

Jean-Philippe Bouchaud (CFM, Ecole Polytechnique)

Alain Comtet (Université P. et M. Curie, Paris VI)

Bertrand Eynard (Commissariat à l'énergie atomique, Saclay)

Jonathan Keating (School of Mathematics, University of Bristol)

Gernot Akemann (Universität Bielefeld)

Aris Moustakas (University of Athens)

Henning Schomerus (Lancaster University)

Vincent Vargas (CNRS, Ecole Normale Supérieure, Paris)

Hans Weidenmüller (Universität Heidelberg)

Anton Zabrodin (ITEP, Moscow)

Random matrix theory and (big) data analysis

Schrödinger operators in random potentials

Topological recursion in random matrices and combinatorics of maps

Random matrix theory and number theory

Matrix models and quantum chromo-dynamics

Random matrix theory methods for telecommunication systems

Random matrix theory approaches to open quantum systems

Gaussian multiplicative chaos and Liouville quantum gravity

Historical overview: random matrix theory and its applications

Some aspects of integrability and quantum-classical correspondence