

# Stability estimates in Sobolev type inequalities

Jean Dolbeault

Ceremade, CNRS & Université Paris-Dauphine  
<http://www.ceremade.dauphine.fr/~dolbeaul>

PDE-Applied Math Seminar

*Department of Mathematics, University of Maryland*

November 21, 2024

# Introduction

• Sobolev inequality on  $\mathbb{R}^d$  with  $d \geq 3$ ,  $2^* = \frac{2d}{d-2}$  and sharp constant  $S_d$

$$\|\nabla f\|_{L^2(\mathbb{R}^d)}^2 \geq S_d \|f\|_{L^{2^*}(\mathbb{R}^d)}^2 \quad \forall f \in \mathcal{D}^{1,2}(\mathbb{R}^d) \quad (\text{S})$$

Equality holds on the manifold  $\mathcal{M}$  of the Aubin–Talenti functions

$$g_{a,b,c}(x) = c (a + |x - b|^2)^{-\frac{d-2}{2}}, \quad a \in (0, \infty), \quad b \in \mathbb{R}^d, \quad c \in \mathbb{R}$$

[Bianchi, Egnell, 1991] there is some non-explicit  $c_{\text{BE}} > 0$  such that

$$\|\nabla f\|_2^2 - S_d \|f\|_{2^*}^2 \geq c_{\text{BE}} \inf_{g \in \mathcal{M}} \|\nabla f - \nabla g\|_2^2$$

• How do we estimate  $c_{\text{BE}}$  ? as  $d \rightarrow +\infty$  ?

Stability & improved entropy – entropy production inequalities

Improved inequalities & faster decay rates for entropies

# Outline

- 1 Explicit stability for Sobolev and LSI on  $\mathbb{R}^d$ 
  - Main results, optimal dimensional dependence; history
  - Sketch of the proof, definitions & preliminary results
  - The main steps of the proof
- 2 Results based on entropy methods and fast diffusion equations
  - Sobolev and HLS inequalities: duality and Yamabe flow
  - Stability, fast diffusion equation and entropy methods
- 3 More stability results for LSI and related inequalities
  - Subcritical interpolation inequalities on the sphere
  - More results on LSI and Gagliardo-Nirenberg inequalities

# Explicit stability results for Sobolev and log-Sobolev inequalities, with optimal dimensional dependence

*Joint papers with M.J. Esteban, A. Figalli, R. Frank, M. Loss*  
**Sharp stability for Sobolev and log-Sobolev inequalities, with  
optimal dimensional dependence**

[arXiv: 2209.08651](#)

**A short review on improvements and stability for some  
interpolation inequalities**

[arXiv: 2402.08527](#)

# An explicit stability result for the Sobolev inequality

Sobolev inequality on  $\mathbb{R}^d$  with  $d \geq 3$ ,  $2^* = \frac{2d}{d-2}$  and sharp constant  $S_d$

$$\|\nabla f\|_{L^2(\mathbb{R}^d)}^2 \geq S_d \|f\|_{L^{2^*}(\mathbb{R}^d)}^2 \quad \forall f \in \dot{H}^1(\mathbb{R}^d) = \mathcal{D}^{1,2}(\mathbb{R}^d)$$

with equality on the manifold  $\mathcal{M}$  of the Aubin–Talenti functions

$$g_{a,b,c}(x) = c (a + |x - b|^2)^{-\frac{d-2}{2}}, \quad a \in (0, \infty), \quad b \in \mathbb{R}^d, \quad c \in \mathbb{R}$$

## Theorem (JD, Esteban, Figalli, Frank, Loss)

*There is a constant  $\beta > 0$  with an explicit lower estimate which does not depend on  $d$  such that for all  $d \geq 3$  and all  $f \in H^1(\mathbb{R}^d) \setminus \mathcal{M}$  we have*

$$\|\nabla f\|_{L^2(\mathbb{R}^d)}^2 - S_d \|f\|_{L^{2^*}(\mathbb{R}^d)}^2 \geq \frac{\beta}{d} \inf_{g \in \mathcal{M}} \|\nabla f - \nabla g\|_{L^2(\mathbb{R}^d)}^2$$

- No compactness argument
- The (estimate of the) constant  $\beta$  is explicit
- The decay rate  $\beta/d$  is optimal as  $d \rightarrow +\infty$

# A stability result for the logarithmic Sobolev inequality

- Use the inverse stereographic projection to rewrite the result on  $\mathbb{S}^d$

$$\begin{aligned} & \|\nabla F\|_{L^2(\mathbb{S}^d)}^2 - \frac{1}{4} d(d-2) \left( \|F\|_{L^{2^*}(\mathbb{S}^d)}^2 - \|F\|_{L^2(\mathbb{S}^d)}^2 \right) \\ & \geq \frac{\beta}{d} \inf_{G \in \mathcal{M}(\mathbb{S}^d)} \left( \|\nabla F - \nabla G\|_{L^2(\mathbb{S}^d)}^2 + \frac{1}{4} d(d-2) \|F - G\|_{L^2(\mathbb{S}^d)}^2 \right) \end{aligned}$$

- Rescale by  $\sqrt{d}$ , consider a function depending only on  $n$  coordinates and take the limit as  $d \rightarrow +\infty$  to approximate the Gaussian measure  $d\gamma = e^{-\pi|x|^2} dx$

## Corollary (JD, Esteban, Figalli, Frank, Loss)

With  $\beta > 0$  as in the result for the Sobolev inequality

$$\begin{aligned} \|\nabla u\|_{L^2(\mathbb{R}^n, d\gamma)}^2 - \pi \int_{\mathbb{R}^n} u^2 \log \left( \frac{|u|^2}{\|u\|_{L^2(\mathbb{R}^n, d\gamma)}^2} \right) d\gamma \\ \geq \frac{\beta \pi}{2} \inf_{a \in \mathbb{R}^n, c \in \mathbb{R}} \int_{\mathbb{R}^n} |u - c e^{a \cdot x}|^2 d\gamma \end{aligned}$$

# Stability for the Sobolev inequality: the history

▷ [Rodemich, 1969], [Aubin, 1976], [Talenti, 1976]

In the inequality  $\|\nabla f\|_{L^2(\mathbb{R}^d)}^2 \geq S_d \|f\|_{L^{2^*}(\mathbb{R}^d)}^2$ , the optimal constant is

$$S_d = \frac{1}{4} d(d-2) |\mathbb{S}^d|^{1-2/d}$$

with equality on the manifold  $\mathcal{M} = \{g_{a,b,c}\}$  of the *Aubin-Talenti functions*

▷ [Lions] a qualitative stability result

$$\text{if } \lim_{n \rightarrow \infty} \|\nabla f_n\|_2^2 / \|f_n\|_{2^*}^2 = S_d, \text{ then } \lim_{n \rightarrow \infty} \inf_{g \in \mathcal{M}} \|\nabla f_n - \nabla g\|_2^2 / \|\nabla f_n\|_2^2 = 0$$

▷ [Brezis, Lieb, 1985] a quantitative stability result ?

▷ [Bianchi, Egnell, 1991] there is some non-explicit  $c_{BE} > 0$  such that

$$\|\nabla f\|_2^2 \geq S_d \|f\|_{2^*}^2 + c_{BE} \inf_{g \in \mathcal{M}} \|\nabla f - \nabla g\|_2^2$$

● The strategy of Bianchi & Egnell involves two steps:

– a local (spectral) analysis: the *neighbourhood* of  $\mathcal{M}$

– a local-to-global extension based on concentration-compactness :

● The constant  $c_{BE}$  is not explicit the *far away regime*

# Stability for the logarithmic Sobolev inequality

- ▷ [Gross, 1975] *Gaussian logarithmic Sobolev inequality* for  $n \geq 1$

$$\|\nabla u\|_{L^2(\mathbb{R}^n, d\gamma)}^2 \geq \pi \int_{\mathbb{R}^n} u^2 \log \left( \frac{|u|^2}{\|u\|_{L^2(\mathbb{R}^n, d\gamma)}^2} \right) d\gamma$$

- ▷ [Weissler, 1979] scale invariant (but dimension-dependent) version of the Euclidean form of the inequality

- ▷ [Stam, 1959], [Federbush, 69], [Costa, 85] Cf. [Villani, 08]

- ▷ [Bakry, Emery, 1984], [Carlen, 1991] equality iff

$$u \in \mathcal{M} := \{w_{a,c} : (a, c) \in \mathbb{R}^d \times \mathbb{R}\} \quad \text{where} \quad w_{a,c}(x) = c e^{a \cdot x} \quad \forall x \in \mathbb{R}^n$$

- [Carlen, 1991] reinforcement of the inequality (Wiener transform)

- ▷ [McKean, 1973], [Beckner, 92] (LSI) as a large  $d$  limit of Sobolev

- ▷ [Bobkov, Gozlan, Roberto, Samson, 2014], [Indrei et al., 2014-23]

stability in Wasserstein distance, in  $W^{1,1}$ , etc.

- ▷ [JD, Toscani, 2016] Comparison with Weissler's form, a (dimension dependent) improved inequality

- ▷ [Fathi, Indrei, Ledoux, 2016] improved inequality assuming a Poincaré inequality (Mehler formula)



# Explicit stability results for the Sobolev inequality

## Proof

# Sketch of the proof

Goal: prove that there is an **explicit** constant  $\beta > 0$  such that for all  $d \geq 3$  and all  $f \in \dot{H}^1(\mathbb{R}^d)$

$$\|\nabla f\|_2^2 \geq S_d \|f\|_{2^*}^2 + \frac{\beta}{d} \inf_{g \in \mathcal{M}} \|\nabla f - \nabla g\|_2^2$$

**Part 1.** We show the inequality for nonnegative functions far from  $\mathcal{M}$   
... *the far away regime*

Make it *constructive*

**Part 2.** We show the inequality for nonnegative functions close to  $\mathcal{M}$   
... *the local problem*

Get *explicit* estimates and remainder terms

**Part 3.** We show that the inequality for nonnegative functions implies the inequality for functions without a sign restriction, up to an acceptable loss in the constant

... *sign-changing functions*

# Some definitions

What we want to minimize is

$$\mathcal{E}(f) := \frac{\|\nabla f\|_{L^2(\mathbb{R}^d)}^2 - S_d \|f\|_{L^{2^*}(\mathbb{R}^d)}^2}{d(f, \mathcal{M})^2} \quad f \in \dot{H}^1(\mathbb{R}^d) \setminus \mathcal{M}$$

where

$$d(f, \mathcal{M})^2 := \inf_{g \in \mathcal{M}} \|\nabla f - \nabla g\|_{L^2(\mathbb{R}^d)}^2$$

▷ up to an elementary *transformation*, we assume that

$d(f, \mathcal{M})^2 = \|\nabla f - \nabla g_*\|_{L^2(\mathbb{R}^d)}^2$  with

$$g_*(x) := |\mathbb{S}^d|^{-\frac{d-2}{2d}} \left( \frac{2}{1+|x|^2} \right)^{\frac{d-2}{2}}$$

▷ use the *inverse stereographic* projection

$$F(\omega) = \frac{f(x)}{g_*(x)} \quad x \in \mathbb{R}^d \text{ with } \begin{cases} \omega_j = \frac{2x_j}{1+|x|^2} & \text{if } 1 \leq j \leq d \\ \omega_{d+1} = \frac{1-|x|^2}{1+|x|^2} \end{cases}$$

# The problem on the unit sphere

Stability inequality on the unit sphere  $\mathbb{S}^d$  for  $F \in H^1(\mathbb{S}^d, d\mu)$

$$\int_{\mathbb{S}^d} (|\nabla F|^2 + A |F|^2) d\mu - A \left( \int_{\mathbb{S}^d} |F|^{2^*} d\mu \right)^{2/2^*} \\ \geq \frac{\beta}{d} \inf_{G \in \mathcal{M}} \left\{ \|\nabla F - \nabla G\|_{L^2(\mathbb{S}^d)}^2 + A \|F - G\|_{L^2(\mathbb{S}^d)}^2 \right\}$$

with  $A = \frac{1}{4} d(d-2)$  and a manifold  $\mathcal{M}$  of optimal functions made of

$$G(\omega) = c (a + b \cdot \omega)^{-\frac{d-2}{2}} \quad \omega \in \mathbb{S}^d \quad (a, b, c) \in (0, +\infty) \times \mathbb{R}^d \times \mathbb{R}$$

- make the reduction of a *far away problem* to a local problem *constructive...* on  $\mathbb{R}^d$
- make the analysis of the *local problem explicit...* on  $\mathbb{S}^d$

## Competing symmetries

• **Rotations on the sphere** combined with stereographic and inverse stereographic projections. Let  $e_d = (0, \dots, 0, 1) \in \mathbb{R}^d$

$$(Uf)(x) := \left( \frac{2}{|x - e_d|^2} \right)^{\frac{d-2}{2}} f \left( \frac{x_1}{|x - e_d|^2}, \dots, \frac{x_{d-1}}{|x - e_d|^2}, \frac{|x|^2 - 1}{|x - e_d|^2} \right)$$

$$\mathcal{E}(Uf) = \mathcal{E}(f)$$

• **Symmetric decreasing rearrangement**  $\mathcal{R}f = f^*$

$f$  and  $f^*$  are equimeasurable

$$\|\nabla f^*\|_{L^2(\mathbb{R}^d)} \leq \|\nabla f\|_{L^2(\mathbb{R}^d)}$$

The method of **competing symmetries**

**Theorem (Carlen, Loss, 1990)**

Let  $f \in L^{2^*}(\mathbb{R}^d)$  be a non-negative function with

$\|f\|_{L^{2^*}(\mathbb{R}^d)} = \|g_*\|_{L^{2^*}(\mathbb{R}^d)}$ . The sequence  $f_n = (\mathcal{R}U)^n f$  is such that

$\lim_{n \rightarrow +\infty} \|f_n - g_*\|_{L^{2^*}(\mathbb{R}^d)} = 0$ . If  $f \in \dot{H}^1(\mathbb{R}^d)$ , then  $(\|\nabla f_n\|_{L^2(\mathbb{R}^d)})_{n \in \mathbb{N}}$  is a non-increasing sequence

## Useful preliminary results

- $\lim_{n \rightarrow \infty} \|f_n - g_*\|_{2^*} = 0$  if  $\|f\|_{2^*} = \|g_*\|_{2^*}$
- $(\|\nabla f_n\|_2^2)_{n \in \mathbb{N}}$  is a nonincreasing sequence

## Lemma

$$\inf_{g \in \mathcal{M}} \|\nabla f - \nabla g\|_2^2 = \|\nabla f\|_2^2 - S_d \sup_{g \in \mathcal{M}, \|g\|_{2^*}=1} (f, g^{2^*-1})^2$$

## Corollary

$(d(f_n, \mathcal{M}))_{n \in \mathbb{N}}$  is strictly decreasing,  $n \mapsto \sup_{g \in \mathcal{M}_1} (f_n, g^{2^*-1})$  is strictly increasing, and

$$\lim_{n \rightarrow \infty} d(f_n, \mathcal{M})^2 = \lim_{n \rightarrow \infty} \|\nabla f_n\|_2^2 - S_d \|g_*\|_{2^*}^2 = \lim_{n \rightarrow \infty} \|\nabla f_n\|_2^2 - S_d \|f\|_{2^*}^2$$

but no monotonicity for  $n \mapsto \mathcal{E}(f_n) = \frac{\|\nabla f_n\|_{L^2(\mathbb{R}^d)}^2 - S_d \|f_n\|_{L^{2^*}(\mathbb{R}^d)}^2}{d(f_n, \mathcal{M})^2}$

# Part 1: Global to local reduction

The *local problem*

$$\mathcal{J}(\delta) := \inf \left\{ \mathcal{E}(f) : f \geq 0, d(f, \mathcal{M})^2 \leq \delta \|\nabla f\|_{L^2(\mathbb{R}^d)}^2 \right\}$$

$f \in \dot{H}^1(\mathbb{R}^d)$  is a nonnegative function in the *far away regime* iff

$$d(f, \mathcal{M})^2 = \inf_{g \in \mathcal{M}} \|\nabla f - \nabla g\|_{L^2(\mathbb{R}^d)}^2 > \delta \|\nabla f\|_{L^2(\mathbb{R}^d)}^2$$

for some  $\delta \in (0, 1)$

Let  $f_n = (\mathcal{R}U)^n f$ . There are two cases:

- (Case 1)  $d(f_n, \mathcal{M})^2 \geq \delta \|\nabla f_n\|_{L^2(\mathbb{R}^d)}^2$  for all  $n \in \mathbb{N}$
- (Case 2) for some  $n \in \mathbb{N}$ ,  $d(f_n, \mathcal{M})^2 < \delta \|\nabla f_n\|_{L^2(\mathbb{R}^d)}^2$

# Global to local reduction – Case 1

$f \in \dot{H}^1(\mathbb{R}^d)$  is a nonnegative function in the far away regime

$$\inf_{g \in \mathcal{M}} \|\nabla f - \nabla g\|_{L^2(\mathbb{R}^d)}^2 > \delta \|\nabla f\|_{L^2(\mathbb{R}^d)}^2$$

## Lemma

Let  $f_n = (\mathcal{R}U)^n f$  and  $\delta \in (0, 1)$

If  $d(f_n, \mathcal{M})^2 \geq \delta \|\nabla f_n\|_{L^2(\mathbb{R}^d)}^2$  for all  $n \in \mathbb{N}$ , then  $\mathcal{E}(f) \geq \delta$

$$\lim_{n \rightarrow +\infty} \|\nabla f_n\|_2^2 \leq \frac{1}{\delta} \lim_{n \rightarrow +\infty} \inf_{g \in \mathcal{M}} \|\nabla f_n - \nabla g\|_2^2 = \frac{1}{\delta} \left( \lim_{n \rightarrow +\infty} \|\nabla f_n\|_2^2 - S_d \|f\|_{2^*}^2 \right)$$

$$1 - \frac{S_d \|f\|_{2^*}^2}{\lim_{n \rightarrow +\infty} \|\nabla f_n\|_2^2} \geq \delta$$

$$\mathcal{E}(f) = \frac{\|\nabla f\|_2^2 - S_d \|f\|_{2^*}^2}{\inf_{g \in \mathcal{M}} \|\nabla f - \nabla g\|_2^2} \geq \frac{\|\nabla f\|_2^2 - S_d \|f\|_{2^*}^2}{\|\nabla f\|_2^2} \geq \frac{\|\nabla f_n\|_2^2 - S_d \|f\|_{2^*}^2}{\|\nabla f_n\|_2^2} \underset{n \rightarrow +\infty}{\geq} \delta$$



## Global to local reduction – Case 2

$$\mathcal{J}(\delta) := \inf \left\{ \mathcal{E}(f) : f \geq 0, d(f, \mathcal{M})^2 \leq \delta \|\nabla f\|_{L^2(\mathbb{R}^d)}^2 \right\} > 0 \text{ (to be proven)}$$

## Lemma

$$\mathcal{E}(f) \geq \delta \mathcal{J}(\delta)$$

$$\text{if } \inf_{g \in \mathcal{M}} \|\nabla f_{n_0} - \nabla g\|_{L^2(\mathbb{R}^d)}^2 > \delta \|\nabla f_{n_0}\|_{L^2(\mathbb{R}^d)}^2$$

$$\text{and } \inf_{g \in \mathcal{M}} \|\nabla f_{n_0+1} - \nabla g\|_{L^2(\mathbb{R}^d)}^2 \leq \delta \|\nabla f_{n_0+1}\|_{L^2(\mathbb{R}^d)}^2$$

▷ Adapt a strategy due to Christ: build a (semi-)continuous rearrangement flow  $(f_\tau)_{n_0 \leq \tau < n_0+1}$  with  $f_{n_0} = Uf_n$  such that  $\|f_\tau\|_{2^*} = \|f\|_2$ ,  $\tau \mapsto \|\nabla f_\tau\|_2$  is nonincreasing, and  $\lim_{\tau \rightarrow n_0+1} f_\tau = f_{n_0+1}$

$$\mathcal{E}(f) \geq 1 - S_d \frac{\|f\|_{2^*}^2}{\|\nabla f\|_2^2} \geq 1 - S_d \frac{\|f_{\tau_0}\|_{2^*}^2}{\|\nabla f_{\tau_0}\|_2^2} = \delta \mathcal{E}(f_{\tau_0}) \geq \delta \mathcal{J}(\delta)$$

Altogether:  $\text{if } d(f, \mathcal{M})^2 > \delta \|\nabla f\|_{L^2(\mathbb{R}^d)}^2, \text{ then } \mathcal{E}(f) \geq \min \{ \delta, \delta \mathcal{J}(\delta) \}$

## Part 2: The (simple) Taylor expansion

### Proposition

Let  $(X, d\mu)$  be a measure space and  $u, r \in L^q(X, d\mu)$  for some  $q \geq 2$  with  $u \geq 0$ ,  $u + r \geq 0$  and  $\int_X u^{q-1} r d\mu = 0$

▷ If  $q = 6$ , then

$$\|u + r\|_q^2 \leq \|u\|_q^2 + \|u\|_q^{2-q} \left( 5 \int_X u^{q-2} r^2 d\mu + \frac{20}{3} \int_X u^{q-3} r^3 d\mu \right. \\ \left. + 5 \int_X u^{q-4} r^4 d\mu + 2 \int_X u^{q-5} r^5 d\mu + \frac{1}{3} \int_X r^6 d\mu \right)$$

▷ If  $3 \leq q \leq 4$ , then

$$\|u + r\|_q^2 - \|u\|_q^2 \\ \leq \|u\|_q^{2-q} \left( (q-1) \int_X u^{q-2} r^2 d\mu + \frac{(q-1)(q-2)}{3} \int_X u^{q-3} r^3 d\mu + \frac{2}{q} \int_X |r|^q d\mu \right)$$

▷ If  $2 < q \leq 3$  (take  $q = 2^*$ ,  $d \geq 6$ ), then

$$\|u + r\|_q^2 \leq \|u\|_q^2 + \|u\|_q^{2-q} \left( (q-1) \int_X u^{q-2} r^2 d\mu + \frac{2}{q} \int_X r_+^q d\mu \right)$$

## Corollary

For all  $\nu > 0$  and for all  $r \in H^1(\mathbb{S}^d)$  satisfying  $r \geq -1$ ,

$$\left(\int_{\mathbb{S}^d} |r|^q d\mu\right)^{2/q} \leq \nu^2 \quad \text{and} \quad \int_{\mathbb{S}^d} r d\mu = 0 = \int_{\mathbb{S}^d} \omega_j r d\mu \quad \forall j = 1, \dots, d+1$$

if  $d\mu$  is the uniform probability measure on  $\mathbb{S}^d$ , then

$$\begin{aligned} \int_{\mathbb{S}^d} (|\nabla r|^2 + A(1+r)^2) d\mu - A \left(\int_{\mathbb{S}^d} (1+r)^q d\mu\right)^{2/q} \\ \geq m(\nu) \int_{\mathbb{S}^d} (|\nabla r|^2 + A r^2) d\mu \end{aligned}$$

$$m(\nu) := \frac{4}{d+4} - \frac{2}{q} \nu^{q-2} \quad \text{if } d \geq 6$$

$$m(\nu) := \frac{4}{d+4} - \frac{1}{3} (q-1)(q-2) \nu - \frac{2}{q} \nu^{q-2} \quad \text{if } d = 4, 5$$

$$m(\nu) := \frac{4}{7} - \frac{20}{3} \nu - 5\nu^2 - 2\nu^3 - \frac{1}{3} \nu^4 \quad \text{if } d = 3$$

An explicit expression of  $\mathcal{J}(\delta)$  if  $\nu > 0$  is small enough so that  $m(\nu) > 0$

## Part 3: Removing the positivity assumption

Take  $f = f_+ - f_-$  with  $\|f\|_{L^{2^*}(\mathbb{R}^d)} = 1$

$$\|\nabla f_+\|_{L^{2^*}(\mathbb{R}^d)}^2 + \|\nabla f_-\|_{L^{2^*}(\mathbb{R}^d)}^2 = \|\nabla f\|_{L^{2^*}(\mathbb{R}^d)}^2$$

Let  $m := \|f_-\|_{L^{2^*}(\mathbb{R}^d)}^{2^*}$  and  $1 - m = \|f_+\|_{L^{2^*}(\mathbb{R}^d)}^{2^*} > 1/2$

$$\begin{aligned} \|f_+\|_{L^{2^*}(\mathbb{R}^d)}^2 + \|f_-\|_{L^{2^*}(\mathbb{R}^d)}^2 - \|f\|_{L^{2^*}(\mathbb{R}^d)}^2 \\ = h_d(m) := m^{\frac{d-2}{d}} + (1-m)^{\frac{d-2}{d}} - 1 \end{aligned}$$

where  $h_d$  is a positive concave function

▷ For some  $g_+ \in \mathcal{M}$

$$\|\nabla f\|_{L^2(\mathbb{R}^d)}^2 - S_d \|f\|_{L^{2^*}(\mathbb{R}^d)}^2 \geq C_{\text{BE}}^d \|\nabla f_+ - \nabla g_+\|_{L^2(\mathbb{R}^d)}^2 + \frac{2h_d(1/2)}{h_d(1/2)+1} \|\nabla f_-\|_{L^2(\mathbb{R}^d)}^2$$

$$C_{\text{BE}}^d \geq \frac{1}{2} \min \left\{ \max_{0 < \delta < 1/2} \delta \mathcal{I}(\delta), \frac{2h_d(1/2)}{h_d(1/2)+1} \right\}$$

## Part 2, refined: The (complicated) Taylor expansion

To get a dimensionally sharp estimate, we expand  $(1+r)^{2^*} - 1 - 2^*r$  with an accurate remainder term for all  $r \geq -1$

$$r_1 := \min\{r, \gamma\}, \quad r_2 := \min\{(r - \gamma)_+, M - \gamma\} \quad \text{and} \quad r_3 := (r - M)_+$$


with  $0 < \gamma < M$ . Let  $\theta = 4/(d-2)$

### Lemma

Given  $d \geq 6$ ,  $r \in [-1, \infty)$ , and  $\bar{M} \in [\sqrt{e}, +\infty)$ , we have

$$\begin{aligned} (1+r)^{2^*} - 1 - 2^*r &\leq \frac{1}{2} 2^* (2^* - 1) (r_1 + r_2)^2 + 2 (r_1 + r_2) r_3 + \left(1 + C_M \theta \bar{M}^{-1} \ln \bar{M}\right) r_3^{2^*} \\ &\quad + \left(\frac{3}{2} \gamma \theta r_1^2 + C_{M, \bar{M}} \theta r_2^2\right) \mathbb{1}_{\{r \leq M\}} + C_{M, \bar{M}} \theta M^2 \mathbb{1}_{\{r > M\}} \end{aligned}$$

where all the constants in the above inequality are explicit



There are constants  $\epsilon_1$ ,  $\epsilon_2$ ,  $k_0$ , and  $\epsilon_0 \in (0, 1/\theta)$ , such that

$$\begin{aligned} \|\nabla r\|_{L^2(\mathbb{S}^d)}^2 + A \|r\|_{L^2(\mathbb{S}^d)}^2 - A \|1 + r\|_{L^{2^*}(\mathbb{S}^d)}^2 \\ \geq \frac{4\epsilon_0}{d-2} \left( \|\nabla r\|_{L^2(\mathbb{S}^d)}^2 + A \|r\|_{L^2(\mathbb{S}^d)}^2 \right) + \sum_{k=1}^3 I_k \end{aligned}$$

$$I_1 := (1 - \theta \epsilon_0) \int_{\mathbb{S}^d} (|\nabla r_1|^2 + A r_1^2) d\mu - A (2^* - 1 + \epsilon_1 \theta) \int_{\mathbb{S}^d} r_1^2 d\mu + A k_0 \theta \int_{\mathbb{S}^d} (r_2^2 \dots)$$

$$I_2 := (1 - \theta \epsilon_0) \int_{\mathbb{S}^d} (|\nabla r_2|^2 + A r_2^2) d\mu - A (2^* - 1 + (k_0 + C_{\epsilon_1, \epsilon_2}) \theta) \int_{\mathbb{S}^d} r_2^2 d\mu$$

$$I_3 := (1 - \theta \epsilon_0) \int_{\mathbb{S}^d} (|\nabla r_3|^2 + A r_3^2) d\mu - \frac{2^*}{2^*} A (1 + \epsilon_2 \theta) \int_{\mathbb{S}^d} r_3^{2^*} d\mu - A k_0 \theta \int_{\mathbb{S}^d} r_3^2 d\mu$$

• spectral gap estimates :  $I_1 \geq 0$

• Sobolev inequality :  $I_3 \geq 0$

• improved spectral gap inequality using that  $\mu(\{r_2 > 0\})$  is small:  $I_2 \geq 0$

[Duoandikoetxea]

# Results based on entropy methods and fast diffusion equations



## Sobolev and Hardy-Littlewood-Sobolev inequalities

- ▷ Stability in a weaker norm, with explicit constants
- ▷ From duality to improved estimates
- ▷ Fast diffusion equation with Yamabe's exponent
- ▷ Explicit stability constants

*Joint paper with G. Jankowiak*  
***Sobolev and Hardy-Littlewood-Sobolev inequalities***  
J. Differential Equations, 257, 2014

# Sobolev and HLS

As it has been noticed by E. Lieb, Sobolev's inequality in  $\mathbb{R}^d$ ,  $d \geq 3$ ,

$$\|\nabla f\|_{L^2(\mathbb{R}^d)}^2 \geq S_d \|f\|_{L^{2^*}(\mathbb{R}^d)}^2 \quad \forall f \in \dot{H}^1(\mathbb{R}^d) = \mathcal{D}^{1,2}(\mathbb{R}^d) \quad (S)$$

and the Hardy-Littlewood-Sobolev inequality

$$\|g\|_{L^{\frac{2d}{d+2}}(\mathbb{R}^d)}^2 \geq S_d \int_{\mathbb{R}^d} g (-\Delta)^{-1} g \, dx \quad \forall g \in L^{\frac{2d}{d+2}}(\mathbb{R}^d) \quad (HLS)$$

are **dual** of each other. Here  $S_d$  is the Aubin-Talenti constant,  $2^* = \frac{2d}{d-2}$ ,  $(2^*)' = \frac{2d}{d+2}$  and by the Legendre transform

$$\sup_{f \in \mathcal{D}^{1,2}(\mathbb{R}^d)} \left( \int_{\mathbb{R}^d} f g \, dx - \frac{1}{2} \|f\|_{L^{2^*}(\mathbb{R}^d)}^2 \right) = \frac{1}{2} \|g\|_{L^{\frac{2d}{d+2}}(\mathbb{R}^d)}^2$$

$$\sup_{f \in \mathcal{D}^{1,2}(\mathbb{R}^d)} \left( \int_{\mathbb{R}^d} f g \, dx - \frac{1}{2} \|\nabla f\|_{L^2(\mathbb{R}^d)}^2 \right) = \frac{1}{2} \int_{\mathbb{R}^d} g (-\Delta)^{-1} g \, dx$$

## Improved Sobolev inequality by duality

## Theorem

[JD, Jankowiak] Assume that  $d \geq 3$  and let  $q = \frac{d+2}{d-2}$

There exists a positive constant  $C \in [\frac{d}{d+4}, 1)$  such that

$$\begin{aligned} \|f^q\|_{L^{\frac{2d}{d+2}}(\mathbb{R}^d)}^2 - S_d \int_{\mathbb{R}^d} f^q (-\Delta)^{-1} f^q dx \\ \leq C S_d^{-1} \|f\|_{L^{2^*}(\mathbb{R}^d)}^{\frac{8}{d-2}} \left( \|\nabla f\|_{L^2(\mathbb{R}^d)}^2 - S_d \|f\|_{L^{2^*}(\mathbb{R}^d)}^2 \right) \end{aligned}$$

for any  $f \in \mathcal{D}^{1,2}(\mathbb{R}^d)$

$C = 1$ : “completion” of the square

$$0 \leq \int_{\mathbb{R}^d} \left| \|f\|_{L^{2^*}(\mathbb{R}^d)}^{\frac{4}{d-2}} \nabla f - S_d \nabla (-\Delta)^{-1} f \right|^2 dx$$

# Using a nonlinear flow to relate Sobolev and HLS

Consider the *fast diffusion* equation

$$\frac{\partial v}{\partial t} = \Delta v^m, \quad t > 0, \quad x \in \mathbb{R}^d \quad (\text{Y})$$

Choice  $m = \frac{d-2}{d+2}$  (Yamabe flow):  $m + 1 = \frac{2d}{d+2}$

## Proposition

Assume that  $d \geq 3$  and  $m = \frac{d-2}{d+2}$ . If  $u = v^m$  and  $v$  is a solution of (Y) with nonnegative initial datum in  $L^{2d/(d+2)}(\mathbb{R}^d)$ , then

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left( S_d^{-1} \|v\|_{L^{\frac{2d}{d+2}}(\mathbb{R}^d)}^2 - \int_{\mathbb{R}^d} v (-\Delta)^{-1} v \, dx \right) \\ &= - \left( \int_{\mathbb{R}^d} v^{m+1} \, dx \right)^{\frac{2}{d}} \left( S_d^{-1} \|\nabla u\|_{L^2(\mathbb{R}^d)}^2 - \|u\|_{L^{2^*}(\mathbb{R}^d)}^2 \right) \leq 0 \end{aligned}$$

Solutions with *separation of variables*

Consider the solution of  $\frac{\partial v}{\partial t} = \Delta v^m$  vanishing at  $t = T$ :

$$\bar{v}_T(t, x) = c (T - t)^\alpha (F(x))^{\frac{d+2}{d-2}}$$

where  $F$  is the Aubin-Talenti solution of

$$-\Delta F = d(d-2) F^{(d+2)/(d-2)}$$

## Lemma

[del Pino, Saez] For any solution  $v$  with initial datum  $v_0 \in L^{2d/(d+2)}(\mathbb{R}^d)$ ,  $v_0 > 0$ , there exists  $T > 0$ ,  $\lambda > 0$  and  $x_0 \in \mathbb{R}^d$  such that

$$\lim_{t \rightarrow T^-} (T - t)^{-\frac{1}{1-m}} \sup_{x \in \mathbb{R}^d} (1 + |x|^2)^{d+2} \left| \frac{v(t, x)}{\bar{v}(t, x)} - 1 \right| = 0$$

with  $\bar{v}(t, x) = \lambda^{(d+2)/2} \bar{v}_T(t, (x - x_0)/\lambda)$

# A convexity improvement

$$J[v] := \int_{\mathbb{R}^d} v^{\frac{2d}{d+2}} dx \quad \text{and} \quad H[v] := S_d \|v\|_{L^{\frac{2d}{d+2}}(\mathbb{R}^d)}^2 - \int_{\mathbb{R}^d} v (-\Delta)^{-1} v dx$$

## Theorem

[JD, Jankowiak] Assume that  $d \geq 3$ . Then we have

$$0 \leq H[v] + S_d J[v]^{1+\frac{2}{d}} \varphi \left( J[v]^{\frac{2}{d}-1} \left( S_d^{-1} \|\nabla u\|_{L^2(\mathbb{R}^d)}^2 - \|u\|_{L^{2^*}(\mathbb{R}^d)}^2 \right) \right)$$

where  $\varphi(x) := \sqrt{1+2x} - 1$  for any  $x \geq 0$

Proof: with  $\kappa_0 := -H'_0/J_0$  and  $H = Y(J)$ , consider the differential inequality

$$Y' \left( C S_d s^{1+\frac{2}{d}} + Y \right) \leq \frac{d+2}{2d} C \kappa_0 S_d^2 s^{1+\frac{4}{d}}, \quad Y(0) = 0, \quad Y(J_0) = H_0$$

# Constructive stability results in Gagliardo-Nirenberg-Sobolev inequalities

*Joint papers with M. Bonforte, B. Nazaret and N. Simonov*  
***Stability in Gagliardo-Nirenberg-Sobolev inequalities: Flows,  
regularity and the entropy method***  
[arXiv:2007.03674](https://arxiv.org/abs/2007.03674), to appear in *Memoirs of the AMS*

***Constructive stability results in interpolation inequalities  
and explicit improvements of decay rates of fast diffusion  
equations***

DCDS, 43 (3&4): 10701089, 2023



# Entropy – entropy production inequality

The fast diffusion equation on  $\mathbb{R}^d$  in self-similar variables

$$\frac{\partial v}{\partial t} + \nabla \cdot \left[ v (\nabla v^{m-1} - 2x) \right] = 0 \quad (\text{FDE})$$

admits a stationary Barenblatt solution  $\mathcal{B}(x) := (1 + |x|^2)^{\frac{1}{m-1}}$

$$\frac{d}{dt} \mathcal{F}[v(t, \cdot)] = -\mathcal{I}[v(t, \cdot)]$$

*Generalized entropy (free energy) and Fisher information*

$$\mathcal{F}[v] := -\frac{1}{m} \int_{\mathbb{R}^d} (v^m - \mathcal{B}^m - m\mathcal{B}^{m-1}(v - \mathcal{B})) \, dx$$

$$\mathcal{I}[v] := \int_{\mathbb{R}^d} v |\nabla v^{m-1} - \nabla \mathcal{B}^{m-1}|^2 \, dx$$

are such that  $\mathcal{I}[v] \geq 4\mathcal{F}[v]$  [del Pino, JD, 2002] so that

$$\mathcal{F}[v(t, \cdot)] \leq \mathcal{F}[v_0] e^{-4t}$$



# Entropy growth rate


$\mathcal{I}[v] \geq 4\mathcal{F}[v] \iff$  Gagliardo-Nirenberg-Sobolev inequalities

$$\|\nabla f\|_{L^2(\mathbb{R}^d)}^\theta \|f\|_{L^{p+1}(\mathbb{R}^d)}^{1-\theta} \geq \mathcal{C}_{\text{GNS}}(p) \|f\|_{L^{2p}(\mathbb{R}^d)} \quad (\text{GNS})$$

with optimal constant. Under appropriate mass normalization

$$v = f^{2p} \text{ so that } v^m = f^{p+1} \text{ and } v |\nabla v^{m-1}|^2 = (p-1)^2 |\nabla f|^2$$

$$p = \frac{1}{2m-1} \iff m = \frac{p+1}{2p} \in [m_1, 1)$$



Asymptotic regime as  $t \rightarrow +\infty$ 

Take  $f_\varepsilon := \mathcal{B}(1 + \varepsilon \mathcal{B}^{1-m} w)$  and expand  $\mathcal{F}[f_\varepsilon]$  and  $\mathcal{I}[f_\varepsilon]$  at order  $O(\varepsilon^2)$   
*linearized free energy and linearized Fisher information*

$$F[w] := \frac{m}{2} \int_{\mathbb{R}^d} w^2 \mathcal{B}^{2-m} dx \quad \text{and} \quad I[w] := m(1-m) \int_{\mathbb{R}^d} |\nabla w|^2 \mathcal{B} dx$$


## Proposition (Hardy-Poincaré inequality)

*[BBDGV, BDNS] Let  $m \in [m_1, 1)$  if  $d \geq 3$ ,  $m \in (1/2, 1)$  if  $d = 2$ , and  $m \in (1/3, 1)$  if  $d = 1$ . If  $w \in L^2(\mathbb{R}^d, \mathcal{B}^{2-m} dx)$  is such that  $\nabla w \in L^2(\mathbb{R}^d, \mathcal{B} dx)$ ,  $\int_{\mathbb{R}^d} w \mathcal{B}^{2-m} dx = 0$ , then*

$$I[w] \geq 4\alpha F[w]$$

*with  $\alpha = 1$ , or  $\alpha = 2 - d(1 - m)$  if  $\int_{\mathbb{R}^d} x w \mathcal{B}^{2-m} dx = 0$*

## Spectral gap



[Denzler, McCann, 2005]

[BBDGV, 2009] [BDGV, 2010] [JD, Toscani, 2010-2015]

Much more is known, *e.g.*, [Denzler, Koch, McCann, 2015]

# The asymptotic time layer improvement

## Proposition

Let  $m \in (m_1, 1)$  if  $d \geq 2$ ,  $m \in (1/3, 1)$  if  $d = 1$ ,  $\eta = 2(dm - d + 1)$  and  $\chi = m/(266 + 56m)$ . If  $\int_{\mathbb{R}^d} v \, dx = \mathcal{M}$ ,  $\int_{\mathbb{R}^d} x v \, dx = 0$  and

$$(1 - \varepsilon) \mathcal{B} \leq v \leq (1 + \varepsilon) \mathcal{B}$$

for some  $\varepsilon \in (0, \chi \eta)$ , then

$$\mathcal{I}[v] \geq (4 + \eta) \mathcal{F}[v]$$

# Uniform convergence in relative error: threshold time

## Theorem

[Bonforte, JD, Nazaret, Simonov, 2021] Assume that  $m \in (m_1, 1)$  if  $d \geq 2$ ,  $m \in (1/3, 1)$  if  $d = 1$  and let  $\varepsilon \in (0, 1/2)$ , small enough,  $A > 0$ , and  $G > 0$  be given. There exists an explicit **threshold time**  $t_* \geq 0$  such that, if  $u$  is a solution of

$$\frac{\partial v}{\partial t} + \nabla \cdot \left[ v (\nabla v^{m-1} - 2x) \right] = 0 \quad (\text{FDE})$$

with nonnegative initial datum  $u_0 \in L^1(\mathbb{R}^d)$  satisfying

$$A[u_0] = \sup_{r>0} r^{\frac{d(m-m_c)}{(1-m)}} \int_{|x|>r} u_0 \, dx \leq A < \infty \quad (\text{H}_A)$$

$\int_{\mathbb{R}^d} u_0 \, dx = \int_{\mathbb{R}^d} B \, dx = \mathcal{M}$ , then

$$\sup_{x \in \mathbb{R}^d} \left| \frac{u(t, x)}{B(t, x)} - 1 \right| \leq \varepsilon \quad \forall t \geq t_*$$

# The initial time layer improvement: backward estimate

By the *carré du champ* method, we have

Away from the Barenblatt solutions,  $\mathcal{Q}[v] := \frac{\mathcal{I}[v]}{\mathcal{F}[v]}$  is such that

$$\frac{d\mathcal{Q}}{dt} \leq \mathcal{Q}(\mathcal{Q} - 4)$$

## Lemma

Assume that  $m > m_1$  and  $v$  is a solution to (FDE) with nonnegative initial datum  $v_0$ . If for some  $\eta > 0$  and  $t_* > 0$ , we have

$\mathcal{Q}[v(t_*, \cdot)] \geq 4 + \eta$ , then


$$\mathcal{Q}[v(t, \cdot)] \geq 4 + \frac{4\eta e^{-4t_*}}{4 + \eta - \eta e^{-4t_*}} \quad \forall t \in [0, t_*]$$

# Stability in Gagliardo-Nirenberg-Sobolev inequalities

*Our strategy*

Choose  $\varepsilon > 0$ , small enough

Get a threshold time  $t_\star(\varepsilon)$



## Two consequences (subcritical case)

▷ Improved decay rate for the fast diffusion equation in rescaled variables

### Corollary

Let  $m \in (m_1, 1)$  if  $d \geq 2$ ,  $m \in (1/2, 1)$  if  $d = 1$ ,  $A > 0$  and  $G > 0$ . If  $v$  is a solution of (FDE) with nonnegative initial datum  $v_0 \in L^1(\mathbb{R}^d)$  such that  $\mathcal{F}[v_0] = G$ ,  $\int_{\mathbb{R}^d} v_0 dx = \mathcal{M}$ ,  $\int_{\mathbb{R}^d} x v_0 dx = 0$  and  $v_0$  satisfies  $(H_A)$ , then

$$\mathcal{F}[v(t, \cdot)] \leq \mathcal{F}[v_0] e^{-(4+\zeta)t} \quad \forall t \geq 0$$

▷ The *stability of the entropy - entropy production inequality*  $\mathcal{I}[v] - 4\mathcal{F}[v] \geq \zeta\mathcal{F}[v]$  also holds in a stronger sense

$$\mathcal{I}[v] - 4\mathcal{F}[v] \geq \frac{\zeta}{4 + \zeta} \mathcal{I}[v]$$



## A constructive stability result (critical case)

Let  $2p^* = 2d/(d-2) = 2^*$ ,  $d \geq 3$  and

$$\mathcal{W}_{p^*}(\mathbb{R}^d) = \left\{ f \in L^{p^*+1}(\mathbb{R}^d) : \nabla f \in L^2(\mathbb{R}^d), |x| f^{p^*} \in L^2(\mathbb{R}^d) \right\}$$

Deficit of the Sobolev inequality:  $\delta[f] := \|\nabla f\|_{L^2(\mathbb{R}^d)}^2 - S_d^2 \|f\|_{L^{2^*}(\mathbb{R}^d)}^2$

### Theorem

Let  $d \geq 3$  and  $A > 0$ . Then for any nonnegative  $f \in \mathcal{W}_{p^*}(\mathbb{R}^d)$  such that

$$\int_{\mathbb{R}^d} (1, x, |x|^2) f^{2^*} dx = \int_{\mathbb{R}^d} (1, x, |x|^2) g dx \quad \text{and} \quad \sup_{r>0} r^d \int_{|x|>r} f^{2^*} dx \leq A$$

we have

$$\|\nabla f\|_{L^2(\mathbb{R}^d)}^2 - S_d^2 \|f\|_{L^{2^*}(\mathbb{R}^d)}^2 \geq \frac{C_*(A)}{4 + C_*(A)} \int_{\mathbb{R}^d} \left| \nabla f + \frac{d-2}{2} f^{\frac{d}{d-2}} \nabla g^{-\frac{2}{d-2}} \right|^2 dx$$

$$C_*(A) = C_*(0) (1 + A^{1/(2d)})^{-1} \quad \text{and} \quad C_*(0) > 0 \text{ depends only on } d$$

# More explicit stability results for the logarithmic Sobolev and Gagliardo-Nirenberg inequalities on $\mathbb{S}^d$

*Joint work with G. Brigati and N. Simonov*

*Logarithmic Sobolev and interpolation inequalities on the sphere: constructive stability results*

Annales IHP, Analyse non linéaire, 362, 2023

*On Gaussian interpolation inequalities*

C. R. Math. Acad. Sci. Paris 41, 2024

# Subcritical interpolation inequalities on the sphere

## ● *Gagliardo-Nirenberg-Sobolev inequality*

$$\|\nabla F\|_{L^2(\mathbb{S}^d)}^2 \geq d \mathcal{E}_p[F] := \frac{d}{p-2} \left( \|F\|_{L^p(\mathbb{S}^d)}^2 - \|F\|_{L^2(\mathbb{S}^d)}^2 \right)$$

for any  $p \in [1, 2) \cup (2, 2^*)$

with  $2^* := \frac{2d}{d-2}$  if  $d \geq 3$  and  $2^* = +\infty$  if  $d = 1$  or  $2$

## ● Limit $p \rightarrow 2$ : the *logarithmic Sobolev inequality*

$$\int_{\mathbb{S}^d} |\nabla F|^2 d\mu \geq \frac{d}{2} \int_{\mathbb{S}^d} F^2 \log \left( \frac{F^2}{\|F\|_{L^2(\mathbb{S}^d)}^2} \right) d\mu \quad \forall F \in H^1(\mathbb{S}^d, d\mu)$$

[Bakry, Emery, 1984], [Bidaud-Véron, Véron, 1991], [Beckner, 1993]

# Gagliardo-Nirenberg inequalities: stability

An improved inequality under orthogonality constraint and the stability inequality arising from the *carré du champ* method can be combined *in the subcritical case* as follows

## Theorem

Let  $d \geq 1$  and  $p \in (1, 2^*)$ . For any  $F \in H^1(\mathbb{S}^d, d\mu)$ , we have

$$\int_{\mathbb{S}^d} |\nabla F|^2 d\mu - d \mathcal{E}_p[F] \geq \mathcal{S}_{d,p} \left( \frac{\|\nabla \Pi_1 F\|_{L^2(\mathbb{S}^d)}^4}{\|\nabla F\|_{L^2(\mathbb{S}^d)}^2 + \|F\|_{L^2(\mathbb{S}^d)}^2} + \|\nabla(\text{Id} - \Pi_1) F\|_{L^2(\mathbb{S}^d)}^2 \right)$$

for some explicit stability constant  $\mathcal{S}_{d,p} > 0$


▷ The result holds true for the logarithmic Sobolev inequality ( $p = 2$ ), again with an explicit constant  $\mathcal{S}_{d,2}$ , for any finite dimension  $d$

Carré du champ – admissible parameters on  $\mathbb{S}^d$ 

[JD, Esteban, Kowalczyk, Loss] Monotonicity of the deficit along

$$\frac{\partial u}{\partial t} = u^{-p(1-m)} \left( \Delta u + (mp - 1) \frac{|\nabla u|^2}{u} \right)$$

$$m_{\pm}(d, p) := \frac{1}{(d+2)^p} \left( dp + 2 \pm \sqrt{d(p-1)(2d - (d-2)p)} \right)$$



**Figure:** Case  $d = 5$ : admissible parameters  $1 \leq p \leq 2^* = 10/3$  and  $m$  (horizontal axis:  $p$ , vertical axis:  $m$ ). Improved inequalities inside !

# Large dimensional limit

*Gagliardo-Nirenberg-Sobolev inequalities on  $\mathbb{S}^d$ ,  $p \in [1, 2)$*

$$\|\nabla u\|_{L^2(\mathbb{S}^d, d\mu_d)}^2 \geq \frac{d}{p-2} \left( \|u\|_{L^p(\mathbb{S}^d, d\mu_d)}^2 - \|u\|_{L^2(\mathbb{S}^d, d\mu_d)}^2 \right)$$

## Theorem

Let  $v \in H^1(\mathbb{R}^n, dx)$  with compact support,  $d \geq n$  and

$$u_d(\omega) = v\left(\omega_1/r_d, \omega_2/r_d, \dots, \omega_n/r_d\right), \quad r_d = \sqrt{\frac{d}{2\pi}}$$

where  $\omega \in \mathbb{S}^d \subset \mathbb{R}^{d+1}$ . With  $d\gamma(y) := (2\pi)^{-n/2} e^{-\frac{1}{2}|y|^2} dy$ ,

$$\begin{aligned} \lim_{d \rightarrow +\infty} d \left( \|\nabla u_d\|_{L^2(\mathbb{S}^d, d\mu_d)}^2 - \frac{d}{2-p} \left( \|u_d\|_{L^2(\mathbb{S}^d, d\mu_d)}^2 - \|u_d\|_{L^p(\mathbb{S}^d, d\mu_d)}^2 \right) \right) \\ = \|\nabla v\|_{L^2(\mathbb{R}^n, d\gamma)}^2 - \frac{1}{2-p} \left( \|v\|_{L^2(\mathbb{R}^n, d\gamma)}^2 - \|v\|_{L^p(\mathbb{R}^n, d\gamma)}^2 \right) \end{aligned}$$

# $L^2$ stability of LSI: comments

[JD, Esteban, Figalli, Frank, Loss]

$$\begin{aligned} \|\nabla u\|_{L^2(\mathbb{R}^n, d\gamma)}^2 - \pi \int_{\mathbb{R}^n} u^2 \log \left( \frac{|u|^2}{\|u\|_{L^2(\mathbb{R}^n, d\gamma)}^2} \right) d\gamma \\ \geq \frac{\beta \pi}{2} \inf_{a \in \mathbb{R}^d, c \in \mathbb{R}} \int_{\mathbb{R}^n} |u - c e^{a \cdot x}|^2 d\gamma \end{aligned}$$

- One dimension is lost (for the manifold of invariant functions) in the limiting process
- Euclidean forms of the stability
- The  $\dot{H}^1(\mathbb{R}^n)$  does not appear, it gets lost in the limit  $d \rightarrow +\infty$
- $\int_{\mathbb{R}^n} |\nabla(u - c e^{a \cdot x})|^2 d\gamma$  ? False, but makes sense under additional assumptions. Some results based on the Ornstein-Uhlenbeck flow and entropy methods: [Fathi, Indrei, Ledoux, 2016], [JD, Brigati, Simonov]
- Taking the limit is difficult because of the lack of compactness

# More results on logarithmic Sobolev inequalities

*Joint work with G. Brigati and N. Simonov*  
***Stability for the logarithmic Sobolev inequality***

Journal of Functional Analysis, 287, oct. 2024

▷ *Entropy methods, with constraints*



# Stability under a constraint on the second moment

$u_\varepsilon(x) = 1 + \varepsilon x$  in the limit as  $\varepsilon \rightarrow 0$

$$d(u_\varepsilon, 1)^2 = \|u'_\varepsilon\|_{L^2(\mathbb{R}, d\gamma)}^2 = \varepsilon^2 \quad \text{and} \quad \inf_{w \in \mathcal{M}} d(u_\varepsilon, w)^\alpha \leq \frac{1}{2} \varepsilon^4 + O(\varepsilon^6)$$

$\mathcal{M} := \{w_{a,c} : (a, c) \in \mathbb{R}^d \times \mathbb{R}\}$  where  $w_{a,c}(x) = c e^{-a \cdot x}$

## Proposition

For all  $u \in H^1(\mathbb{R}^d, d\gamma)$  such that  $\|u\|_{L^2(\mathbb{R}^d)} = 1$  and  $\|xu\|_{L^2(\mathbb{R}^d)}^2 \leq d$ , we have

$$\|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 - \frac{1}{2} \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma \geq \frac{1}{2d} \left( \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma \right)^2$$

and, with  $\psi(s) := s - \frac{d}{4} \log(1 + \frac{4}{d}s)$ ,

$$\|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 - \frac{1}{2} \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma \geq \psi \left( \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \right)$$

# Stability under log-concavity

## Theorem

For all  $u \in H^1(\mathbb{R}^d, d\gamma)$  such that  $u^2 \gamma$  is log-concave and such that

$$\int_{\mathbb{R}^d} (1, x) |u|^2 d\gamma = (1, 0) \quad \text{and} \quad \int_{\mathbb{R}^d} |x|^2 |u|^2 d\gamma \leq K$$

we have

$$\|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 - \frac{\mathcal{C}_\star}{2} \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma \geq 0$$

$$\mathcal{C}_\star = 1 + \frac{1}{432K} \approx 1 + \frac{0.00231481}{K}$$

Self-improving Poincaré inequality and stability for LSI

[Fathi, Indrei, Ledoux, 2016]

## Theorem

Let  $d \geq 1$ . For any  $\varepsilon > 0$ , there is some explicit  $\mathcal{C} > 1$  depending only on  $\varepsilon$  such that, for any  $u \in H^1(\mathbb{R}^d, d\gamma)$  with

$$\int_{\mathbb{R}^d} (1, x) |u|^2 d\gamma = (1, 0), \quad \int_{\mathbb{R}^d} |u|^2 e^{\varepsilon|x|^2} d\gamma < \infty$$

for some  $\varepsilon > 0$ , then we have

$$\|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \geq \frac{\mathcal{C}}{2} \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma$$

with  $\mathcal{C} = 1 + \frac{\mathcal{C}_*(K_*) - 1}{1 + R^2 \mathcal{C}_*(K_*)}$ ,  $K_* := \max\left(d, \frac{(d+1)R^2}{1+R^2}\right)$  if  $\text{supp}(u) \subset B(0, R)$

Compact support: [Lee, Vázquez, '03]; [Chen, Chewi, Niles-Weed, '21]

These slides can be found at

<http://www.ceremade.dauphine.fr/~dolbeaul/Lectures/>  
▷ Lectures

More related papers can be found at

<http://www.ceremade.dauphine.fr/~dolbeaul/Preprints/list/>  
▷ Preprints and papers

For final versions, use **Dolbeault** as login and **Jean** as password

E-mail: [dolbeaul@ceremade.dauphine.fr](mailto:dolbeaul@ceremade.dauphine.fr)

Thank you for your attention !